

NAME: _____

MARK BOX		
PROBLEM	POINTS	
1	10	
2	10	
3	10	
4	10	
TOTAL	40	

ID (last four digits) _____

please check the box of your section below

or

INSTRUCTIONS:

- (1) To receive credits you must:
- work in a logical fashion, **show all your work and indicate your reasoning** to support and justify your answer
 - when applicable put your answer on/in the line/box; use the back of the page if needed
- (2) This exam covers (from *Elementary Linear Algebra* by Larson 8th ed.):
Sections 4.1–4.5, 4.6*, 7.1 .

- (1) Let $A = \begin{pmatrix} 1 & 9 \\ 0 & -1 \end{pmatrix}$. Verify that (i) the eigenvalues are $\lambda_1 = 1$ and $\lambda_2 = -1$
(ii) the corresponding eigenvectors of the matrix A are given respectively as

$$t \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad t \begin{pmatrix} -9 \\ 2 \end{pmatrix}$$

for all t in $(-\infty, \infty)$.

- (2) Find (i) the characteristic equation, (ii) the eigenvalues, and (iii) the corresponding eigenvectors of the matrix. (iv*) [optional] Is the matrix diagonalizable?
- (a)

$$\begin{pmatrix} 4 & -5 \\ 2 & -3 \end{pmatrix}$$

(b)

$$\begin{pmatrix} 1 & -1 & -1 \\ 1 & 3 & 1 \\ -3 & 1 & -1 \end{pmatrix}$$

- (3) (optional)* Find the adjoint $\mathbf{ad}(M)$ of the matrix $M = \begin{pmatrix} -1 & 0 & 2 \\ 0 & 3 & 2 \\ 3 & 0 & -1 \end{pmatrix}$.

Verify that $M\mathbf{ad}(M) = \mathbf{ad}(M)M = \det(M)I_3$.

- (4) **Definition.** A vector \mathbf{u} is said to be in the null space of a matrix A provided

$$A\mathbf{u} = \mathbf{0}.$$

or, equivalently, \mathbf{u} is an eigenvector corresponding to the zero eigenvalue of A .

Which of the following vectors, if any, is in the null space of $A = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 2 & 1 & 1 & 3 \\ 1 & 0 & 2 & 2 \end{pmatrix}$?

- a) $[-1 \ 0 \ 1 \ 0]^T$ b) $[0 \ 2 \ 1 \ -1]^T$ c) $[0 \ 4 \ 2 \ -2]^T$
- (5) Determine which of the following statements are equivalent to the fact that a matrix A of size $n \times n$ is invertible?
- a) A is nonsingular
 b) The row space of A has dimension n
 c) The column space of A has dimension n
 d) The determinant of A is nonzero
 e) The system $A\mathbf{x} = \mathbf{b}$ has a unique solution for any given \mathbf{b} in \mathbf{R}^n
 f) The system $A\mathbf{x} = \mathbf{0}$ has nonzero solution
 g) The dimension of the null space of A is zero
 h) The rows of A are linear independent
 i) The columns of A are linear independent
 j) The rank of A is n
 k) A is row-equivalent to an identity matrix
 l) All eigenvalues of A are nonzero
 m) A can be written as the product of elementary matrices.
- (6) (optional*) The matrix $A = \begin{pmatrix} 2 & 1 & 3 & 1 \\ 1 & -1 & 0 & 1 \\ 1 & 1 & 2 & 1 \end{pmatrix}$ row reduces to $C = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$.
- a) Find the rank and nullity of A .
 b) Find a basis of the row space and the column space of A respectively.
 c) Find a basis of the null space of A
 d) Does the system $A\mathbf{x} = \begin{pmatrix} 109 \\ -217 \\ 66 \end{pmatrix}$ have a solution? (Hint: You can draw a conclusion from the fact that dimension of column space is 3, without having to solve the system. Recall that $\text{rank}(A) = \dim(\text{Col}(A)) = \dim(\text{Row}(A))$)
 e) What is the relation between $\text{rank}, \dim(\text{null}(A))$? (Hint: Theorem 4.17 (pp.196) states that $\text{rank}(A) + \dim(\text{null}(A)) = n$, the number of columns)
- (7) Find all the eigenvalues of the given matrix.
- a) $\begin{pmatrix} 1 & -2 & 0 \\ -3 & 1 & 0 \\ -4 & -5 & 1 \end{pmatrix}$

b) $\begin{pmatrix} 1 & 9 \\ 0 & -1 \end{pmatrix}$ (c) $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ (d) $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ (e) $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$
 where $i = \sqrt{-1}$ ($i^2 = -1$) is the unit for pure imaginary numbers.

(8) We say a vector \mathbf{u} is a linear combination of a finite set of vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ if there exist constants c_1, c_2, c_3 such that

$$\mathbf{u} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3.$$

Determine whether one can write $\mathbf{u} = [8 \ 3 \ 8]^T$ as a linear combination of the vectors in the set S .

$$S = \{[4 \ 3 \ 2]^T, [0 \ 3 \ 2]^T, [0 \ 0 \ 2]^T\}$$

Solutions 2 (a). (i) The characteristic equation is $|\lambda I - A| = 0$, that is,

$$\begin{vmatrix} \lambda - 4 & 5 \\ -2 & \lambda + 3 \end{vmatrix} = \lambda^2 - \lambda - 2 = (\lambda + 1)(\lambda - 2) = 0$$

(ii) The eigenvalues are solutions to the characteristic equation:

$$\lambda_1 = -1, \lambda_2 = 2.$$

(iii) The eigenvectors corresponding to $\lambda = -1$ is the set of nonzero solutions to $(\lambda I - A)\mathbf{x} = \mathbf{0}$

$$\begin{pmatrix} -5 & 5 \\ -2 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Solving it yields

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad t \neq 0$$

Similarly the eigenvectors corresponding to $\lambda = 2$ are

$$\begin{pmatrix} -2 & 5 \\ -2 & 5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Solving it yields

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = t \begin{pmatrix} 5 \\ 2 \end{pmatrix} \quad t \neq 0$$

2 (b). (i) The characteristic equation reads

$$\begin{vmatrix} \lambda - 1 & 1 & 1 \\ -1 & \lambda - 3 & -1 \\ 3 & -1 & \lambda + 1 \end{vmatrix} = 0$$

(ii) The eigenvalues are obtained by solving the above equation. We start with simplifying

$$\begin{aligned} & \begin{vmatrix} \lambda - 1 & 1 & 1 \\ -1 & \lambda - 3 & -1 \\ 3 & -1 & \lambda + 1 \end{vmatrix} = \begin{vmatrix} \lambda - 2 & \lambda - 2 & 0 \\ -1 & \lambda - 3 & -1 \\ 3 & -1 & \lambda + 1 \end{vmatrix} \\ & = (\lambda - 2) \begin{vmatrix} 1 & 1 & 0 \\ -1 & \lambda - 3 & -1 \\ 3 & -1 & \lambda + 1 \end{vmatrix} = (\lambda - 2) \begin{vmatrix} 1 & 0 & 0 \\ -1 & \lambda - 2 & -1 \\ 3 & -4 & \lambda + 1 \end{vmatrix} \\ & = (\lambda - 2) \begin{vmatrix} \lambda - 2 & -1 \\ -4 & \lambda + 1 \end{vmatrix} \\ & = (\lambda - 2)(\lambda + 2)(\lambda - 3). \end{aligned}$$

Hence $\lambda_1 = -2$, $\lambda_2 = 2$ and $\lambda_3 = 3$.

2 (b) (iii) To find the eigenvectors for λ , we solve the linear homogeneous equation

$$\begin{bmatrix} \lambda - 1 & 1 & 1 \\ -1 & \lambda - 3 & -1 \\ 3 & -1 & \lambda + 1 \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

If $\lambda_1 = -2$, row reduction yields

$$\begin{aligned} & \begin{bmatrix} \lambda_1 - 1 & 1 & 1 \\ -1 & \lambda_1 - 3 & -1 \\ 3 & -1 & \lambda_1 + 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -\frac{1}{4} \\ 0 & 1 & \frac{1}{4} \\ 0 & 0 & 0 \end{bmatrix} \\ & \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = t \begin{pmatrix} \frac{1}{4} \\ -\frac{1}{4} \\ 1 \end{pmatrix} \quad t \neq 0. \end{aligned}$$

The eigenvectors for λ_2 and λ_3 can be found in a similar way. If $\lambda_3 = 3$, say, row reduction yields

$$\begin{aligned} & \begin{bmatrix} \lambda_3 - 1 & 1 & 1 \\ -1 & \lambda_3 - 3 & -1 \\ 3 & -1 & \lambda_3 + 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix} \\ & \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = t \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \quad t \neq 0. \end{aligned}$$

3*. By definition the adjoint matrix of a matrix $A = (C_{ij})_{n \times n}$ is given by

$$\mathbf{ad}(A) = \begin{pmatrix} C_{11} & C_{21} & \cdots & C_{n1} \\ C_{12} & C_{22} & \cdots & C_{n2} \\ C_{1n} & C_{2n} & \cdots & C_{nn} \end{pmatrix}$$

where $C_{ij} = (-1)^{i+j} M_{ij}$ are cofactors of A .

$$\begin{pmatrix} -3 & 0 & -6 \\ 6 & -5 & 2 \\ -9 & 0 & -3 \end{pmatrix}$$

A straight forward computation shows $M\mathbf{ad}(M) = \mathbf{ad}(M)M = -15I_3$.

4. Answer: (b) and (c). If multiplying A and the vector in (b), we will have $A\mathbf{u} = 0$. The same occurs for (c).

(Here is some more details. Given a matrix A , the null space $Null(A)$ is a vector space consisting of all those vectors \mathbf{u} satisfying the equation $A\mathbf{x} = 0$.

So if you want to check if certain vector u is in the null space, all you need to do is to substitute $\mathbf{x} = \mathbf{u}$ into the linear equation $A\mathbf{x} = 0$.

If you find $A\mathbf{u} = 0$ then \mathbf{u} belongs to $Null(A)$; otherwise it does not belong to $Null(A)$.

6*. (a) $Rank(A) = 3$. $nullity(A) = 1$ (nullity is the dimension for the null space of A)

(b) A basis for $Row(A)$ is given by $\{[2 \ 1 \ 3 \ 1]^T, [1 \ -1 \ 0 \ 1]^T, [1 \ 1 \ 2 \ 1]^T\}$. A basis for $Col(A)$ is given by $\{[2 \ 1 \ 1]^T, [3 \ 0 \ 2]^T, [1 \ 1 \ 1]^T\}$.

(c) The solutions to $A\mathbf{x} = \mathbf{0}$ consist vectors of the form $\{t \begin{pmatrix} -1 \\ -1 \\ 1 \\ 0 \end{pmatrix}, t \neq 0\}$. So a basis can be

chosen as $\begin{pmatrix} -1 \\ -1 \\ 1 \\ 0 \end{pmatrix}$.

(d) Yes. Because the dimension of the column space of A equals to 3, and, the dimension of the column space of the augmented matrix $[Ab]$ is also 3. We see that the column space and the augmented space are consistent in the case. Therefore the system $A\mathbf{x} = \mathbf{b}$ is consistent or solvable.

(e) $Rank(A) + \dim(null(A)) = 3 + 1 = 4$ which should be the number of columns.

7. (a) The eigenvalues are solutions of

$$\begin{vmatrix} \lambda - 1 & 2 & 0 \\ 3 & \lambda - 1 & 0 \\ 4 & 5 & \lambda - 1 \end{vmatrix} = 0$$

$$(\lambda - 1) \begin{vmatrix} \lambda - 1 & 2 \\ 3 & \lambda - 1 \end{vmatrix} = (\lambda - 1)(\lambda^2 - 2\lambda - 5) = 0.$$

Hence $\lambda_1 = 1$, $\lambda_{2,3} = 1 \pm \sqrt{6}$.

7 (c). $\lambda = \pm i$.

7 (d) $\lambda = \pm 1$.

7. (e) Solving

$$\begin{vmatrix} \lambda & -i \\ -i & \lambda \end{vmatrix} = \lambda^2 + 1 = 0$$

we obtain $\lambda_1 = i$, $\lambda_2 = -i$.

(8) We can rewrite $\mathbf{u} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3$ in the form

$$\begin{pmatrix} 4 & 0 & 0 \\ 3 & 3 & 0 \\ 2 & 2 & 2 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 8 \\ 3 \\ 8 \end{pmatrix}.$$

Solve this equation using either row reduction or in the traditional way as follows.

$$\begin{cases} 4c_1 = 8 \\ 3c_1 + 3c_2 = 3 \\ 2c_1 + 2c_2 + 2c_3 = 8 \end{cases} \Rightarrow \begin{cases} c_1 = 2 \\ c_1 + c_2 = 1 \\ c_1 + c_2 + c_3 = 4 \end{cases} \Rightarrow$$

$$\therefore \mathbf{c} = [c_1, c_2, c_3]^T = [2 \ -1 \ 3]^T$$